

## Plant schedule

## 81 Grove Road, Sutton

Item	Spring	Summer	Autumn	Winter
Anemone × hybrida 'Königin Charlotte'	Apply 5 - 7 cm mulch.		Cut back Stems when flowering is over	
Anemone hupehensis Hadspen Abundance	Apply 5 - 7 cm mulch.		Cut back Stems when flowering is over	
Anemone japonicus honorine jobert	Apply 5 - 7 cm mulch.		Cut back Stems when flowering is over	
Campanula portenschlagiana			Cut back after flowering	
			Apply a slow-release balanced fertiliser and a mulch of	
Caryopteris × clandonensis 'Heavenly Blue@	Prune all of the flowered stems back hard to the		well-rotted garden compost around the base of the	
	permanent framework.Apply 5 - 7 cm mulch.		plant	
Erigeron karvinskianus			Trim back lanky stems	
	Apply a slow-release balanced fertiliser and a mulch			
Exochorda x macrantha Niagara	of well-rotted garden compost around the base of			
	the plant			
Committee Bossess		Cut back once flowers have faded to encourage fresh new		
Geranium Rozanne		growth and a second flush of flowers	Cut back once flowering is over in late autumn	
	Apply a light mulch (3cm) of well-rotted garden			
Hakonechloa macra	compost after cutting back old foliage and before			
	new growth emerges			
Helleborus hybridus	Provide a top-dressing of general fertiliser		Cut back the stalks after the flowers have faded	Remove old leaves
	Carefully trim back in April, taking care not to cut into			
Lavandula angustifolia Munstead	old wood.		Once established, trim back lanky stems after flowering	
			and remove up to a third of the oldest stems	
	Apply a generous 5-7cm (2-3in) mulch of well-rotted			
Lonicera Rhubarb and Custard	compost or manure around the base of the plant,			
	avoiding the immediate crown.	Tie in to wires or trellis		
	Lift and divide large clumps in spring, replanting			
Nepeta Walkers Low	divided specimens with lots of well-rotted organic	Cut back once flowers have faded to encourage fresh new		
	matter.	growth and a second flush of flowers		Cut back after it has finished flowering,
	Lift and divide large clumps in spring, replanting			
Persicaria bistorta superba	divided specimens with lots of well-rotted organic			
	matter.	Dead head as flowers fade to encourage fresh new growth	Cut back after flowering	
Philadelphus 'Snowbelle'	Mulch with a thick layer of compost or well rotted		Prune in late summer, removing one in four stems to	
,	manure		ground	
Prunus 'Beni-yutaka'	Mulch with a thick layer of compost or well rotted			
	manure avoiding stem		Check ties and stakes.	
Prunus Pink Perfection	Mulch with a thick layer of compost or well rotted			
	manure avoiding stem		Check ties and stakes.	
	Lightly trim shoots after flowering. Apply a generous			
Sarcococca confusa	5-7cm (2-3in) mulch of well-rotted compost around			
	the base of the plant			
Stachys officinalis 'Hummelo'				
•		Dead head as flowers fade to encourage fresh new growth	Cut back after flowering	
Taxus baccata ball	Mulch with a thick layer of compost or well rotted	<del>-</del>	<del>-</del> :	
	manure	Trim to shape late summer /early autumn	Trim to shape late summer /early autumn	

Trachelospermum jasminoides	Mulch with a thick layer of compost or well rotted		Trim back to fit available space
	manure	Tie in to wires or trellis	Tilli back to lit available space
Weigela 'Bristol Ruby'	Mulch with a thick layer of compost or well rotted manure	Once established, in midsummer after flowering prune back a	
		few of the older stems to base and cut back the shoots to a	
		strong bud below the spent flowers	